

# NAVAL POSTGRADUATE SCHOOL

## Monterey, California



FORTRAN SUBROUTINES FOR THE EVALUATION OF THE  
CONFLUENT HYPERGEOMETRIC FUNCTIONS

WILLIAM GRAGG  
BENY NETA

August 1989

Approved for public release; distribution unlimited  
Prepared for:

Naval Postgraduate School  
Monterey, CA 93943

1-2000  
7-30-00, 1/2, 1/2, 1/2-58-97-14

NAVAL POSTGRADUATE SCHOOL  
Department of Mathematics

Rear Admiral R. W. West, JR.  
Superintendent

Harrison Shull  
Provost

This report was prepared in conjunction with research funded by the Naval Postgraduate School Research Council. Reproduction of all or part of this report is authorized.

Prepared by:

Prepared by:

## REPORT DOCUMENTATION PAGE

OUTLINE OF CHUCK LIBRARY  
NAVAL POSTGRADUATE SCHOOL  
CALIFORNIA 93943 600

1 REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b RESTRICTIVE MARKINGS	
2 SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2 DECLASSIFICATION/DOWNGRADING SCHEDULE			
4 PERFORMING ORGANIZATION REPORT NUMBER(S) NPS-53-89=014		5 MONITORING ORGANIZATION REPORT NUMBER(S) NPS-53-89-014	
6a NAME OF PERFORMING ORGANIZATION Naval Postgraduate School	6b OFFICE SYMBOL (If applicable) 53	7a NAME OF MONITORING ORGANIZATION Naval Postgraduate School	
6c ADDRESS (City, State, and ZIP Code) Monterey, CA 93943		7b ADDRESS (City, State, and ZIP Code) Monterey, CA 93943	
8a NAME OF FUNDING/SPONSORING ORGANIZATION Naval Postgraduate School	8b OFFICE SYMBOL (If applicable) 53	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER O&MN, Direct Funding	
8c ADDRESS (City, State, and ZIP Code) Monterey, CA 93943		10 SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO	PROJECT NO
		TASK NO	WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) FORTRAN SUBROUTINES FOR THE EVALUATION OF THE CONFLUENT HYPERGEOMETRIC FUNCTIONS			
12 PERSONAL AUTHOR(S) William Gragg and Beny Neta			
13a TYPE OF REPORT Technical Report	13b TIME COVERED FROM 2/89 TO 8/89	14 DATE OF REPORT (Year, Month, Day) 89 August 14	15 PAGE COUNT 12
16 SUPPLEMENTARY NOTATION			
COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		confluent hypergeometric functions, stable algorithm Fortran subroutine, recurrence relation	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) <p>In this report we list the Fortran subroutines for evaluating the confluent hypergeometric functions <math>M(a,b;x)</math> and <math>U(a,b;x)</math>. These subroutines use the stable recurrence relations given e.g. in Wimp.</p>			
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a NAME OF RESPONSIBLE INDIVIDUAL William Gragg and Beny Neta		22b TELEPHONE (Include Area Code) (408) 646-2194, 2235	22c OFFICE SYMBOL 53Gr and 53Nd

FORM 1473, 84 MAR

83 APR edition may be used until exhausted  
All other editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE  
UNCLASSIFIED



Fortran Subroutines for the Evaluation of the  
Confluent Hypergeometric Functions

W. Gragg  
and  
B. Neta

Naval Postgraduate School  
Department of Mathematics  
Monterey, CA 93943



## Abstract

In this report we list the Fortran subroutines for evaluating the confluent hypergeometric functions  $M(a,b;x)$  and  $U(a,b;x)$ . These subroutines use the stable recurrence relations given e.g. in Wimp.

Key words:  
confluent hypergeometric functions  
stable algorithm  
Fortran subroutine  
recurrence relation

## Introduction

It is well known that the ordinary differential equation

$$x \frac{d^2 y}{dx^2} + (1-x) \frac{dy}{dx} - ay = 0$$

has a solution

$$y(x) = AM(a,1;x) + BU(a,1;x)$$

if  $a$  is not a negative integer.

This problem arises e.g. when solving the linearized shallow water equations with the full linear variation in depth included (see Williams, Staniforth and Neta, [1]).

The computation of the confluent hypergeometric functions is based on the Miller algorithm (see e.g. Wimp, [2]). In general, one has a second order difference equation

$$z(n) + a(n)z(n+1) + b(n)z(n+2) = 0, \quad n \geq 0, \quad b(n) \neq 0.$$

If  $b(n) = 0$  for some  $n$ , in some cases one can make a change of variable  $Y(n) = \lambda(n)z(n)$  which will produce an equation of the desired type. Let  $w(n)$  be a nontrivial solution and the sum of the normalizing series

$$S = \sum_{k=0}^{\infty} c(k)w(k) \neq 0$$



is known. Let  $N$  be a large integer and define  $z_N(n)$ ,  $0 \leq n \leq N+1$ , by

$$z_N(n) = \begin{cases} 0 & n = N+1 \\ 1 & n = N \end{cases}$$

$$z_N(n) + a(n)z_N(n+1) + b(n)z_N(n+2) = 0, \quad n = N-1, \dots, 1, 0.$$

One can approximate  $w(n)$  by  $w_N(n)$

$$w_N(n) = Sz_N(n)/S_N$$

where

$$S_N = \sum_{k=0}^N c(k)z_N(k).$$

The algorithm is said to converge if

$$w_N(n) \rightarrow w(n) \quad \text{as} \quad N \rightarrow \infty.$$

The function  $M(a,b;x)$  satisfies the recurrence relation

$$\begin{aligned} (2n+b+2)(n+a)z(n) - (2n+b+1)\left\{(2a-b) + \frac{(2n+b)(2n+b+2)}{x}\right\}z(n+1) \\ - (2n+b)(n+b+1-a)z(n+2) = 0. \end{aligned}$$

The minimal solution is

$$w(n) = \frac{x^n (a)_n}{(b)_{2n}} M(a+n, 2n+b; x)$$

where

$$(c)_n = \frac{\Gamma(n+c)}{\Gamma(c)} .$$

The normalization relationship used in our subroutine is

$$S = b-1 = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} (b-1)_k (b+2k-1) w(k) .$$

An obvious modification must be made if  $b = 1$ . The algorithm is not defined if  $b$ ,  $b+1-a$ ,  $a$  are negative integers or zero.

The function  $U(a,b;x)$  satisfies the relationship

$$\begin{aligned} (n+a)(n+a+1-b)z(n) - (n+1)[2(n+a+1)+x-b]z(n+1) \\ + (n+1)(n+2)z(n+2) = 0 . \end{aligned}$$

The minimal solution is

$$w(n) = \frac{x^n (a)_n (a+1-b)_n}{n!} U(a+n, b; x)$$

for  $|\arg x| < \pi$ . A normalization relation is

$$1 = \sum_{k=0}^{\infty} w(k) .$$

In the next section we give a listing of the Fortran subroutines.

# Subroutine Miller

```
SUBROUTINE MILLER(N,ALPHA,BETA,X,S,SS,COEFF)
INTEGER N
REAL*8 ALPHA,BETA,X,SS
REAL*8 S(0:1000)
EXTERNAL COEFF
C USES THE J.C.P. MILLER ALGORITHM TO COMPUTE
C S(0:N).
C BEGIN
    INTEGER NN,K
    REAL*8 T,D,EPS,A,B,C
    REAL*8 OLDS(0:1000)
    EPS = 0.000000001
C    INITIALIZE OLDS.
    DO 1 K = 0, 1000
        OLDS(K) = 0
1    CONTINUE
C    CHOOSE INITIAL NN.
    NN = N + 2
C    INITIALIZE K, S AND T.
    2    K = NN
        S(K+1) = 0
        S(K) = 1
        CALL COEFF(K,ALPHA,BETA,X,A,B,C)
        T = 2*C*S(K)
C    TAKE A BACKWARD RECURRENCE STEP AND UPDATE IT.
    3    K = K - 1
        CALL COEFF(K,ALPHA,BETA,X,A,B,C)
        S(K) = A*S(K+1) + B*S(K+2)
C    CHECK FOR OVERFLOW AND RESCALE IF NECESSARY.
        D= DABS(S(K))
        IF (D .GT. 1.D30) THEN
C        BEGIN
            CALL SCALE(K,NN,S,T,D)
        END IF
        IF (K .GT. 0) THEN
C        BEGIN
            T = T + 2*C*S(K)
            GO TO 3
        END IF
        T = T + C*S(0)
        DO 4 K = 0, N
            S(K) = S(K)/T
4    CONTINUE
C    TEMPORARY PRINT STATEMENT.
C    PRINT*, S(0)
C    TEST FOR CONVERGENCE.
        D = 0
        DO 5 K = 0, N
            D = D + S(K)**2
5    CONTINUE
        D = DSQRT(D)
        T = 0
```

```

      DO 6 K = 0, N
        T = T + (S(K) - OLDS(K))**2
6      CONTINUE
      T = DSQRT(T)
C      TAKE ANOTHER STEP IF NO CONVERGENCE.
      IF (T .GT. EPS*D) THEN
C      BEGIN
        NN = 2*NN
        DO 7 K = 0, N
          OLDS(K) = S(K)
7        CONTINUE
        IF(NN .LE. 1000) GO TO 2
        PRINT 999,NN,ALPHA,BETA,X,T
999      FORMAT(' ** NO CONVERGENCE ** NN AP CP X T ',I5,4E14.7)
      END IF
      SS=S(0)
      RETURN
END

```

```

SUBROUTINE COEFF(N,ALPHA,BETA,X,A,B,C)
INTEGER N
REAL*8 ALPHA,BETA,X,A,B,C
C COMPUTES COEFFICIENTS USED BY J.C.P. MILLER ALGORITHM FOR
C A CONFLUENT HYPERGEOMETRIC FUNCTION  $M(a,b;x)$ 
C SEE JET WIMP. COMPUTATION WITH RECURRENCE RELATIONS.
C PITMAN 1984 PP. 61-62
C BEGIN
    INTEGER M,K
    REAL*8 T,U,V,W
    S = 2*ALPHA - BETA
    T = N + ALPHA
    M = 2*N
    U = M + BETA
    V = U + 1
    W = V + 1
    A = (S/W + U/X)*V/T
    B = (N + BETA - ALPHA + 1)*U/T/W
    T = 1
    IF (N .GT. 0) THEN
C BEGIN
        S = BETA - 1
        DO 1 K = 1, N-1
            T = -T*(1+S/K)
1        CONTINUE
            T = -T*(1+S/M)
        END IF
        C = T
        RETURN
    END
END

```

```

SUBROUTINE SCALE(K,N,S,T,D)
INTEGER N,K
REAL*8 T,D
REAL*8 S(0:1000)
C BEGIN
    INTEGER J
    T = T/D
    DO 1 J = K, N
        S(J) = S(J)/D
1    CONTINUE
    RETURN
END

```

```

SUBROUTINE COEFU(N,ALPHA,BETA,X,A,B,C)
INTEGER N
REAL*8 ALPHA, BETA,X,A,B,C
C COMPUTES COEFFICIENTS USED BY J.C.P. MILLER ALGORITHM FOR
C A CONFLUENT HYPERGEOMETRIC FUNCTION  $U(a,b;x)$ 
C SEE JET WIMP, COMPUTATION WITH RECURRENCE RELATIONS,
C PITMAN 1984 PP. 63-64
C BEGIN
      INTEGER M,K
      REAL*8 S,T,U,V,W
      S = ALPHA + QFLOAT(N)
      T = S + 1.D0
      U = S*(T - BETA)
      V = QFLOAT(N + 1)
      W = V + 1.D0
      A = (2*T + X - BETA)*V/U
      B = - V*W/U
      C = 1
      RETURN
END

```

Remark: The program that calls Miller must supply as a last parameter either COEFF (for M) or COEFU (for U).

The subroutines are available on a diskette from either author upon request. These subroutines were tested extensively for various values of  $a$ ,  $b$  and  $x$ .

Remark: If the parameter is a negative integer, the solution of the differential equation is

$$y = AL_n(x) + B\{\ln|x|L_n(x) + \sum_{m=0}^{\infty} \beta_m x^m\}$$

where  $n = -a$ .

$L_n(x)$  are Laguerre polynomials whose coefficients  $\alpha_i$  satisfy

$$\alpha_i = \frac{i-n-1}{i^2} \alpha_{i-1} \quad i = 2, \dots, n,$$

$$\alpha_1 = -n.$$

The coefficients  $\beta_m$  satisfy

$$\beta_{m+1} = \frac{(m-n)\beta_m + \left(1 - \frac{2(m-n)}{m+1} \alpha_m\right)}{(m+1)^2} \quad m = 1, \dots, n-1$$

$$\beta_m = \frac{1}{(n+1)^2} \alpha_n \quad m = n$$

$$\beta_m = \frac{m-n-1}{m^2} \beta_{m-1} \quad m = n+1, n+2, \dots$$

### Acknowledgement:

This research was conducted for the Office of Naval Research and was funded by the Naval Postgraduate School.

### References

1. R.T. Williams, A.N. Staniforth and B. Neta, Solution of a generalized Sturm-Liouville Problem, IMA Conference on Computational Ordinary Differential Equations, Imperial College, London, July 3-7, 1989.
2. J. Wimp. Computation with Recurrence Relations, Pitman Advanced Pub. Program, Boston, 1984.



# DISTRIBUTION LIST

	NO. OF COPIES
Director Defense Tech. Information Center Cameron Station Alexandria, VA 22314	2
Director of Research Administration Code 012 Naval Postgraduate School Monterey, CA 93943	1
Library Code 0142 Naval Postgraduate School Monterey, CA 93943	2
Department of Mathematics Code 53 Naval Postgraduate School Monterey, CA 93943	1
Center for Naval Analyses 4401 Ford Avenue Alexandria, VA 22302-0268	1
Professor Beny Neta Code 53Nd Department of Mathematics Naval Postgraduate School Monterey, CA 93943	6
Dr. C.P. Katti J. Nehru University School of Computer & Systems Sciences New Delhi 110067 India	5
Professor Paul Nelson Texas A&M University Department of Nuclear Engineering and Mathematics College Station, TX 7 843-3133	1
Professor H.B. Keller Department of Applied Mathematics California Institute of Technology Pasadena, CA 91125	1

Professor W. Gragg Code 53Gr Department of Mathematics Naval Postgraduate School Monterey, CA 93943	6
Profesor H. Dean Victory, Jr. Texas Tech University Department of Mathematics Lubbock, TX 79409	1
Professor Gordon Latta Code 53Lz Department of Mathematics Naval Postgraduate School Monterey, CA 93943	1
Professor Arthur Schoenstadt Code 53Zh Department of Mathematics Naval Postgraduate School Monterey, CA 93943	1
Professor M.M. Chawla, Head Department of Mathematics III/III/B-1, IIT Campus Hauz Khas, New Delhi 110016 India	1
Professor R.T. Williams Code 63Wu Department of Meteorology Naval Postgraduate School Monterey, CA 93943	1



DUDLEY KNOX LIBRARY



3 2768 00333153 9